

## **FACS I Unit 19**

### **Student Supplement 3**

1. Answers should include three of the following: ectopic pregnancy, preterm delivery, stillbirths, birth defects, illness, and death.
2. Most babies become infected during delivery while passing through the birth canal of an infected mother.
3. Babies infected with chlamydia during delivery can develop eye infections and pneumonia.
4. If babies contract gonorrhea at birth, it can affect their eyes and joints and cause severe blood infections.
5. Syphilis can cause fetal or infant death and preterm delivery. Babies can also develop brain damage, blindness, hearing loss, teeth problems, and bone abnormalities.
6. Some babies develop sores on the skin or in the mouth. Sometimes a baby's infection can cause brain damage, blindness, mental retardation, or death.
7. Women should make sure they do not have an STI before becoming pregnant or contract an STI during pregnancy. Sexually active women should be in a monogamous relationship with a partner who is not infected with an STI.

### **Student Supplement 4**

1. Answers should include three of the following: abstinence is 100% effective in preventing the spread of sexually transmitted diseases, there are no side effects, no doctor's visit or prescription is necessary, no cost, lowers the risk of a woman developing cervical cancer, lowers the risk of infertility.
2. Smoking increases the risk of developing blood clots, heart attack, and stroke.
3. Condoms help protect against the spread of sexually transmitted diseases.
4. A woman who misses a day of her birth control pill should take it as soon as she remembers, even if it means taking two in one day. If she misses two pills, she should call her doctor.
5. Sterilization is intended to be permanent. It is not appropriate for anyone who may eventually want to have children. Young men or women who have the procedure may change their mind in the future.